


FILE COPY



PubMed	Nucleotide	Protein	Genome	Structure	Popset
Search	Protein	<input checked="" type="checkbox"/> for		Go	Clear
Limits		Index	History	Clipboard	
Display	Default	<input checked="" type="checkbox"/> View as	HTML	<input checked="" type="checkbox"/> Save	Add to Clipboard
Hide Brief and Lin					

☐ 1: GI = "114939" [GenPept] BETA-GALACTOSIDASE (LACTASE) PubMed, Related Sequences,

LOCUS BGAL\_ECOLI 1024 aa BCT 15-JUL-1999  
DEFINITION BETA-GALACTOSIDASE (LACTASE).  
ACCESSION P00722  
PID g114939  
VERSION P00722 GI:114939  
DBSOURCE swissprot: locus BGAL\_ECOLI, accession P00722;  
class: standard.  
created: Jul 21, 1986.  
sequence updated: Jul 21, 1986.  
annotation updated: Jul 15, 1999.  
xrefs: gi: 146575, gi: 146577, gi: 1786532, gi: 1786539, gi:  
1657477, gi: 1657540, gi: 41901, gi: 1197203, gi: 41897, gi: 41898,  
gi: 67496  
xrefs (non-sequence databases): ECO2DBASE E123.0, ECOGENE EG10527,  
PFAM PF00703, PROSITE PS00719, PROSITE PS00608  
KEYWORDS Hydrolase; Glycosidase; 3D-structure.  
SOURCE Escherichia coli.  
ORGANISM Escherichia coli  
Bacteria; Proteobacteria; gamma subdivision; Enterobacteriaceae;  
Escherichia.  
REFERENCE 1 (residues 1 to 1024)  
AUTHORS Kalnins,A., Otto,K., Ruther,U. and Muller-Hill,B.  
TITLE Sequence of the lacZ gene of Escherichia coli  
JOURNAL EMBO J. 2 (4), 593-597 (1983)  
MEDLINE 84028567  
REMARK SEQUENCE FROM N.A.  
REFERENCE 2 (residues 1 to 1024)  
AUTHORS BLATTNER,F.R., PLUNKETT,G.I.I.I., BLOCH,C.A., PERNA,N.T.,  
BURLAND,V., RILEY,M., COLLADO-VIDES,J., GLASNER,F.D., RODE,C.K.,  
MAYHEW,G.F., GREGOR,J., DAVIS,N.W., KIRKPATRICK,H.A., GOEDEN,M.A.,  
ROSE,D.J., MAU,B. and SHAO,Y.  
TITLE The complete genome sequence of Escherichia coli K-12  
JOURNAL Science 277 (5331), 1453-1474 (1997)  
MEDLINE 97426617  
REMARK SEQUENCE FROM N.A.  
STRAIN=K12 / MG1655  
REFERENCE 3 (residues 1 to 1024)  
AUTHORS DUNCAN,M., ALLEN,E., ARAUJO,R., APARICIO,A.M., CHUNG,E., DAVIS,K.,  
FEDERSPIEL,N., HYMAN,R., KALMAN,S., KOMP,C., KURDI,O., LEW,H.,  
LIN,D., NAMATH,A., OEFNER,P., ROBERTS,D., SCHRAMM,S. and DAVIS,R.W.  
TITLE Direct Submission  
JOURNAL Submitted (??-NOV-1996) to the EMBL/GenBank/DDBJ databases  
REMARK SEQUENCE FROM N.A.  
REFERENCE 4 (residues 1 to 1024)  
AUTHORS Fowler,A.V. and Zabin,I.  
TITLE Amino acid sequence of beta-galactosidase. XI. Peptide ordering  
procedures and the complete sequence

JOURNAL J. Biol. Chem. 253 (15), 5521-5525 (1978)  
 MEDLINE 78218239  
 REMARK SEQUENCE.  
 REFERENCE 5 (residues 1 to 1024)  
 AUTHORS Calos, M.P. and Miller, J.H.  
 TITLE Molecular consequences of deletion formation mediated by the  
 transposon Tn9  
 JOURNAL Nature 285 (5759), 38-41 (1980)  
 MEDLINE 80188189  
 REMARK SEQUENCE OF 355-475 FROM N.A.  
 REFERENCE 6 (residues 1 to 1024)  
 AUTHORS Buchel, D.E., Gronenborn, B. and Muller-Hill, B.  
 TITLE Sequence of the lactose permease gene  
 JOURNAL Nature 283 (5747), 541-545 (1980)  
 MEDLINE 80120651  
 REMARK SEQUENCE OF 1007-1023 FROM N.A.  
 REFERENCE 7 (residues 1 to 1024)  
 AUTHORS Fowler, A.V. and Smith, P.J.  
 TITLE The active site regions of lacZ and ebg beta-galactosidases are  
 homologous  
 JOURNAL J. Biol. Chem. 258 (17), 10204-10207 (1983)  
 MEDLINE 83290932  
 REMARK ACTIVE SITE REGIONS.  
 REFERENCE 8 (residues 1 to 1024)  
 AUTHORS Herrchen, M. and Legler, G.  
 TITLE Identification of an essential carboxylate group at the active site  
 of lacZ beta-galactosidase from Escherichia coli  
 JOURNAL Eur. J. Biochem. 138 (3), 527-531 (1984)  
 MEDLINE 84108409  
 REMARK ACTIVE SITE GLU-461.  
 REFERENCE 9 (residues 1 to 1024)  
 AUTHORS Gebler, J.C., Aebersold, R. and Withers, S.G.  
 TITLE Glu-537, not Glu-461, is the nucleophile in the active site of (lac  
 Z) beta-galactosidase from Escherichia coli  
 JOURNAL J. Biol. Chem. 267 (16), 11126-11130 (1992)  
 MEDLINE 92283812  
 REMARK ACTIVE SITE GLU-537.  
 REFERENCE 10 (residues 1 to 1024)  
 AUTHORS Jacobson, R.H., Zhang, X.J., DuBose, R.F. and Matthews, B.W.  
 TITLE Three-dimensional structure of beta-galactosidase from E. coli  
 JOURNAL Nature 369 (6483), 761-766 (1994)  
 MEDLINE 94277211  
 REMARK X-RAY CRYSTALLOGRAPHY (3.5 ANGSTROMS).  
 COMMENT

-----  
 This SWISS-PROT entry is copyright. It is produced through a  
 collaboration between the Swiss Institute of Bioinformatics and  
 the EMBL outstation - the European Bioinformatics Institute.  
 The original entry is available from <http://www.expasy.ch/sprot>  
 and <http://www.ebi.ac.uk/sprot>  
 -----

[CATALYTIC ACTIVITY] HYDROLYSIS OF TERMINAL, NON-REDUCING  
 BETA-D-GALACTOSE RESIDUES IN BETA-D-GALACTOSIDES.  
 [SUBUNIT] HOMOTETRAMER.  
 [SIMILARITY] BELONGS TO FAMILY 2 OF GLYCOSYL HYDROLASES.  
 [DATABASE] NAME=Worthington enzyme manual;  
 WWW='http://www.worthington-biochem.com/manual/G/BG.html'.  
 [DATABASE] NAME=ProZyme technical fact sheet;  
 WWW='http://www.prozyme.com/technical/bgalddata.html'.  
 -----

FEATURES  
 source Location/Qualifiers  
 1..1024  
 /organism="Escherichia coli"  
 /db\_xref="taxon:562"  
 1..1024  
 Protein 1..1024  
 /product="BETA-GALACTOSIDASE"

Site /EC\_number="3.2.1.23"  
462  
/site\_type="active"  
/note="PROTON DONOR."  
Site 538  
/site\_type="active"  
/note="NUCLEOPHILE."

## ORIGIN

```
1 mtmitdslav vlqrrdwenp gvtqlnrlaa hppfaswrns eeartdrpsq qlrslngewr
61 fawfpapeav peswlecdlp eadtvvvpsn wqmhgdydapi ytnvtypitv nppfvptenp
121 tgcysltfnv deswlqegqt riifdgvnas fhlcwngwv gygqdsrlps efdlsaflra
181 genrlavmvl rwsdgsyled qdmwrmsgif rdvsllhkpt tqisdfhvat rfnddfsrav
241 leaevqmcge lrdylrvtvs lwqgetqvas gtapfggeii derggyadv trlrnvenpk
301 lwsaeipnly ravvelhtad gtlieaeacd vgfrevrien gllllngkpl lirgvnrheh
361 hplhgqvmde qtmvqdillm kqnnfnavr shypnhplwy tldryglyv vdeaniethg
421 mvpmnrltdd prwlpamser vtrmvqrdrn hpsviiwslg nesghganhd alyrwiksvd
481 psrpvqyegg gadttatdii cpmyarvded qpfpavpkws ikkwlsilpge trplilceya
541 hamgnslggf akywqafqy prlqggfvwd wvdqslkyd engnpwsayg gdfgdtndr
601 qfcmnglvfa drtphpalte akhqgqffqf rlsqgtievt seylfrhsdn ellhwmvald
661 gkplasevpl ldvapqgkql ielpelpqpe sagqlwltvr vvgpnataws eaghisaawq
721 wrlaenlsvt lpaashaiph lttsemdfci elgnkrwqfn rqsqflsqmw igdkkqltpt
781 lrdqftrapl dndigvseat ridpnawver wkaaghyqae aallqctadt ladavlitta
841 hawqhggktl fisrkyrid gsgqmaitvd vevsdtphp ariglncqla qvaervnwlg
901 lgpqenydr ltaacfdwrw lplsdmytpy vfpsenglrc gtrelnygph qwrgdfqfni
961 srysqqqlme tshrhllhae egtwnidgf hmgiggddsw spsvsaefql sagryhyqlv
1021 wcqk
```

//

Restrictions on Use | Write to the HelpDesk  
NCBI | NLM | NIH

L12 ANSWER 1 OF 6 BIOSIS COPYRIGHT 2000 BIOSIS  
AN 2000:148109 BIOSIS  
DN PREV2000000148109  
TI Isolation of neuronal precursor cells by **FACS** from the Talphal-  
**GFP transgenic** animals.  
AU Sawamoto, K. (1); Ogawa, Y. (1); Saiwaki, T. (1); Kawaguchi, A. (1);  
Miyata, T. (1); Goldman, S. A.; Okano, H.  
CS (1) Dept. of Neuroanatomy, Biomed. Res. Ctr., Osaka Univ. Med. Sch.,  
Suita, Osaka, 565-0871 Japan  
SO Society for Neuroscience Abstracts., (1999) Vol. 25, No. 1-2, pp. 2043.  
Meeting Info.: 29th Annual Meeting of the Society for Neuroscience. Miami  
Beach, Florida, USA October 23-28, 1999 Society for Neuroscience  
. ISSN: 0190-5295.  
DT Conference  
LA English  
SL English

R(321.N48)

L2 ANSWER 9 OF 10 BIOSIS COPYRIGHT 2000 BIOSIS  
AN 1998:528162 BIOSIS  
DN PREV199800528162  
TI Visualization of neuronal stem cells in vivo.  
AU Yamaguchi, Masahiro (1); Suzuki, Misao; Mori, Kensaku  
CS (1) PRESTO, Japan Sci. Technol. Corp., Soraku-gu, Kyoto 619-0237 Japan  
SO Neuroscience Research Supplement, (1998) No. 22, pp. S286.  
Meeting Info.: 21st Annual Meeting of the Japan Neuroscience Society and  
the First Joint Meeting of the Japan Neuroscience Society and the Japanese  
Society for Neurochemistry Tokyo, Japan September 21-23, 1998 Japan  
Neuroscience Society  
. ISSN: 0921-8696.  
DT Conference  
LA English

L8 ANSWER 2 OF 2 MEDLINE  
 AN 97388554 MEDLINE  
 DN 97388554  
 TI Brief expression of a GFP cre fusion gene in embryonic stem cells allows rapid retrieval of site-specific genomic deletions.  
 AU Gagnetten S; Le Y; Miller J; Sauer B  
 CS Laboratory of Biochemistry and Metabolism, National Institute of Diabetes, Digestive and Kidney Disease, National Institutes of Health, Bethesda, MD 20892-1800, USA.  
 SO NUCLEIC ACIDS RESEARCH, (1997 Aug 15) 25 (16) 3326-31.  
 Journal code: O8L. ISSN: 0305-1048.  
 CY ENGLAND: United Kingdom  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals; Cancer Journals  
 EM 199711  
 EW 19971103  
 AB The Cre DNA recombinase of bacteriophage P1 has become a useful tool for precise genomic manipulation in embryonic stem (ES) cells that have been gene modified by homologous recombination. We have re-engineered the cre gene to allow ready identification of living Cre+cells by constructing a functional fusion between Cre and an enhanced **green** fluorescent protein from Aequorea victoria (GFPS65T). The GFP cre fusion gene product rapidly targeted the nucleus in the absence of any exogenous nuclear localization signal. Moreover, GFPCre catalyzed efficient DNA recombination in both a mouse 3T3 derivative cell line and in murine ES cells. Fluorescence- activated cell sorting (**FACS**) of transiently GFP cre -transfected ES cells not only allowed rapid and efficient isolation of Cre+cells after DNA transfection but also demonstrated that a burst of Cre expression is sufficient to commit cells to Cre-mediated 'pop-out' of loxP -tagged DNA from the genome. Thus, GFP cre allows rapid identification of living cells in which loxP - flanked DNA sequences are destined to be removed from the genome by Cre-mediated recombination without reliance on recombinational activation or inactivation of a marker gene at the target locus. In addition, the GFP cre fusion gene will prove useful in tracing tissue-specific Cre expression in **transgenic** animals, thereby facilitating the generation and analysis of conditional gene knockout mice.

41, 24, 13

L2 ANSWER 36 OF 37 MEDLINE  
 AN 94121904 MEDLINE  
 DN 94121904  
 TI Independent regulatory elements in the **nestin** gene direct  
**transgene** expression to neural stem cells or muscle precursors  
 [published erratum appears in Neuron 1994 Jun;12(6):following 1388].  
 AU Zimmerman L; Parr B; Lendahl U; Cunningham M; McKay R; Gavin B; Mann J;  
 Vassileva G; McMahon A  
 CS Department of Brain and Cognitive Sciences, Massachusetts Institute of  
 Technology, Cambridge 02139..  
 SO NEURON, (1994 Jan) 12 (1) 11-24. *qp 356.2 N48*  
 Journal code: AN8. ISSN: 0896-6273.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199405  
 AB Changes in intermediate filament gene expression occur at key steps in the  
 differentiation of cell types in the mammalian CNS. Neuroepithelial stem  
 cells express the intermediate filament protein **nestin** and  
 down-regulate it sharply at the transition from proliferating stem cell to  
 postmitotic neuron. **Nestin** is also expressed in muscle  
 precursors but not in mature muscle cells. We show here that in  
**transgenic** mice, independent cell type-specific elements in the  
 first and second introns of the **nestin** gene consistently direct  
 reporter gene expression to developing muscle and neural precursors,  
 respectively. The second intron contains an enhancer that functions in CNS  
 stem cells, suggesting that there may be a single transcriptional  
 mechanism regulating the CNS stem cell state. This enhancer is much less  
 active in the PNS. The identification of these elements facilitates  
 analysis of mechanisms controlling the switch in gene expression that  
 occurs when muscle and brain precursors terminally differentiate.

what promoter??  
 Nestin Rat

L2 ANSWER 33 OF 37 MEDLINE  
 AN 96017714 MEDLINE  
 DN 96017714  
 TI Rapid, widespread, and longlasting induction of **nestin** contributes to the generation of glial scar tissue after CNS injury.  
 AU Frisen J; Johansson C B; Torok C; Risling M; Lendahl U  
 CS Department of Neuroscience, Karolinska Institute, Stockholm, Sweden..  
 SO JOURNAL OF CELL BIOLOGY, (1995 Oct) 131 (2) 453-64.  
 Journal code: HMM. ISSN: 0021-9525.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals; Cancer Journals  
 EM 199602  
 AB Neuronal regeneration does generally not occur in the central nervous system (CNS) after injury, which has been attributed to the generation of glial scar tissue. In this report we show that the composition of the glial scar after traumatic CNS injury in rat and mouse is more complex than previously assumed: expression of the intermediate filament **nestin** is induced in reactive astrocytes. **Nestin** induction occurs within 48 hours in the spinal cord both at the site of lesion and in degenerating tracts and lasts for at least 13 months. **Nestin** expression is induced with similar kinetics in the crushed optic nerve. In addition to the expression in reactive astrocytes, we also observed **nestin** induction within 48 hours after injury in cells close to the central canal in the spinal cord, while **nestin** expressing cells at later timepoints were found progressively further out from the central canal. This dynamic pattern of **nestin** induction after injury was mimicked by lacZ expressing cells in **nestin** promoter/lacZ **transgenic** mice, suggesting that defined **nestin** regulatory regions mediate the injury response. We discuss the possibility that the spatiotemporal pattern of **nestin** expression reflects a population of **nestin** positive cells, which proliferates and migrates from a region close to the central canal to the site of lesion in response to injury.

11 Enhancer?



L2 ANSWER 30 OF 37 MEDLINE  
 AN 97147786 MEDLINE  
 DN 97147786  
 TI Bypass of lethality with mosaic mice generated by Cre-loxP-mediated recombination.  
 AU Betz U A; Vosschenrich C A; Rajewsky K; Muller W  
 CS Institute for Genetics, University of Cologne, Germany..  
 ubetz@mac.genetik.uni-koeln.de  
 SO CURRENT BIOLOGY, (1996 Oct 1) 6 (10) 1307-16.  
 Journal code: B44. ISSN: 0960-9822.  
 CY ENGLAND: United Kingdom  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199704  
 EW 19970403  
 AB BACKGROUND: The analysis of gene function based on the generation of mutant mice by homologous recombination in embryonic stem cells is limited if gene disruption results in embryonic lethality. Mosaic mice, which contain a certain proportion of mutant cells in all organs, allow lethality to be circumvented and the potential of mutant cells to contribute to different cell lineages to be analyzed. To generate mosaic animals, we used the bacteriophage P1-derived Cre-loxP recombination system, which allows gene alteration by Cre-mediated deletion of loxP-flanked gene segments. RESULTS: We generated **nestin-cre transgenic** mouse lines, which expressed the Cre recombinase under the control of the rat **nestin** promoter and its second intron enhancer. In crosses to animals carrying a loxP-flanked target gene, partial deletion of the loxP-flanked allele occurred before day 10.5 post coitum and was detectable in all adult organs examined, including germ-line cells. Using this approach, we generated mosaic mice containing cells deficient in the gamma-chain of the interleukin-2 receptor (IL-2R gamma); in these animals, the IL-2R gamma-deficient cells were underrepresented in the thymus and spleen. Because mice deficient in DNA polymerase beta die perinatally, we studied the effects of DNA polymerase beta deficiency in mosaic animals. We found that some of the mosaic polymerase beta-deficient animals were viable, but were often reduced in size and weight. The fraction of DNA polymerase beta-deficient cells in mosaic embryos decreased during embryonic development, presumably because wild-type cells had a competitive advantage. CONCLUSIONS: The **nestin-cre transgenic** mice can be used to generate mosaic animals in which target genes are mutated by Cre-mediated recombination of loxP-flanked target genes. By using mosaic animals, embryonic lethality can be bypassed and cell lineages for whose development a given target gene is critical can be identified. In the case of DNA polymerase beta, deficient cells are already selected against during embryonic development, demonstrating the general importance of this protein in multiple cell types.

is cre fluorescent?

L2 ANSWER 29 OF 37 MEDLINE  
 AN 97258113 MEDLINE  
 DN 97258113  
 TI An evolutionarily conserved region in the second intron of the human **nestin** gene directs gene expression to CNS progenitor cells and to early neural crest cells.  
 AU Lothian C; Lendahl U  
 CS Department of Cell and Molecular Biology, Medical Nobel Institute, Karolinska Institute, Stockholm, Sweden.  
 SO EUROPEAN JOURNAL OF NEUROSCIENCE, (1997 Mar) 9 (3) 452-62.  
 Journal code: BYG. ISSN: 0953-816X.  
 CY ENGLAND: United Kingdom  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199709  
 EW 19970901  
 AB Central nervous system (CNS) progenitor cells transiently proliferate in the embryonic neural tube and give rise to neurons and glial cells. A characteristic feature of the CNS progenitor cells is expression of the intermediate filament **nestin** and it was previously shown that the rat **nestin** second intron functions as an enhancer, directing gene expression to CNS progenitor cells. In this report we characterize the **nestin** enhancer in further detail. Cloning and sequence analysis of the rat and human **nestin** second introns revealed local domains of high sequence similarity in the 3' portion of the introns. **Transgenic** mice were generated with the most conserved 714 bp in the 3' portion of the intron, or with the complete, 1852 bp, human second intron, coupled to the reporter gene lacZ. The two constructs gave a very similar **nestin**-like expression pattern, indicating that the important control elements reside in the 714 bp element. Expression was observed starting in embryonic day (E)7.5 neural plate, and at E10.5 CNS progenitor cells throughout the neural tube expressed lacZ. At E12.5, lacZ expression was more restricted and confined to proliferating regions in the neural tube. An interesting difference, compared to the rat **nestin** second intron, was that the human intron at E10.5 mediated lacZ expression also in early migrating neural crest cells, which is a site of endogenous **nestin** expression. In conclusion, these data show that a relatively short, evolutionarily conserved region is sufficient to control gene expression in CNS progenitor cells, but that the same region differs between rodents and primates in its capacity to control expression in neural crest cells.

promoter?

L2 ANSWER 28 OF 37 MEDLINE

AN 97359974 MEDLINE

DN 97359974

TI Limb proprioceptive deficits without neuronal loss in **transgenic** mice overexpressing neurotrophin-3 in the developing nervous system.

AU Ringstedt T; Kucera J; Lendahl U; Ernfors P; Ibanez C F

CS Department of Neuroscience, Karolinska Institute, Stockholm, Sweden.

SO DEVELOPMENT, (1997 Jul) 124 (13) 2603-13.

Journal code: ECW. ISSN: 0950-1991.

CY ENGLAND: United Kingdom

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199710

AB The role of neurotrophin-3 (NT3) during sensory neuron development was investigated in **transgenic** mice overexpressing NT3 under the control of the promoter and enhancer regions of the **nestin** gene, an intermediate filament gene widely expressed in the developing nervous system. Most of these mice died during the first postnatal day, and all showed severe limb ataxia suggestive of limb proprioceptive dysfunction. Tracing and histological analyses revealed a complete loss of spindles in limb muscles, absence of peripheral and central Ia projections, and lack of cells immunoreactive to parvalbumin in the dorsal root ganglion (DRG). Despite these deficits, there was no neuronal loss in the DRG of these mice. At birth, **transgenic** DRG showed increased neuron numbers, and displayed a normal proportion of neurons expressing substance P, calcitonin gene-related peptide and the NT3 receptor trkC. **Transgenic** dorsal roots exhibited an increased number of axons at birth, indicating that all sensory neurons in **transgenic** mice projected to the dorsal spinal cord. Despite the absence of central Ia afferents reaching motoneurons, several sensory fibers were seen projecting towards ectopic high levels of NT3 in the midline of **transgenic** spinal cords. These findings suggest novel roles for NT3 in differentiation of proprioceptive neurons, target invasion and formation of Ia projections which are independent from its effects on neuronal survival.

102(b)

NT3 has 4W

11y

out of 258 ans

L2 ANSWER 27 OF 37 MEDLINE

AN 97391813 MEDLINE

DN 97391813

TI **Transgenic** analysis of central nervous system development and regeneration.

AU Lendahl U

CS Department of Cell and Molecular Biology, Medical Nobel Institute, Karolinska Institute, Stockholm, Sweden.

SO ACTA ANAESTHESIOLOGICA SCANDINAVICA. SUPPLEMENTUM, (1997) 110 116-8. Journal code: 08Q. ISSN: 0515-2720.

CY Denmark

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199711

EW 19971103

AB The **transgenic** technique allows specific genetic alterations to be made in all cells of an animal and this has greatly improved our understanding of how the embryonic and adult central nervous system (CNS) develop. The CNS originates from the neuroectoderm in the neural plate on the dorsal side of the embryo and after closure of the neural tube the cells of the neuroepithelium, i.e. the CNS stem cells, transiently proliferate to generate neurons and glial cells. Here we review our attempts to gain insights into the control of CNS development. We have identified a gene, **nestin**, which is predominantly expressed in embryonic and adult CNS stem cells. In addition to its normal expression in the CNS stem cells, **nestin** is reexpressed in CNS tumors and in the adult spinal cord and brain after CNS injury. By using the lacZ reporter gene assay in **transgenic** mice, we have identified regulatory regions (enhancer) in the **nestin** gene required for expression in embryonic CNS stem cells and in the adult spinal cord after injury. In a second project, we have cloned and characterized the Notch gene family (the Notch 1, 2 and 3 genes) in mouse and man. These genes encode trans-membrane receptors, which appear to be key regulatory molecules for proliferation and differentiation both in the developing CNS and in other tissues. Expression of an activated form of the Notch 3 receptor from the **nestin** promoter in **transgenic** mice leads to a lethal, exencephaly-like phenotype in the embryo, probably as a result of excess proliferation of the CNS stem cells. The recent finding that the Notch 3 gene is the genetic cause for familial stroke is discussed in the context of current models for Notch function.

L2 ANSWER 13 OF 37 MEDLINE  
AN 1999117355 MEDLINE  
DN 99117355  
TI Heterogeneity of neural progenitor cells revealed by enhancers in the  
**nestin** gene.  
AU Yaworsky P J; Kappen C  
CS Molecular Neuroscience Program, Samuel C. Johnson Medical Research Center,  
Scottsdale, Arizona, 85259, USA.  
SO DEVELOPMENTAL BIOLOGY, (1999 Jan 15) 205 (2) 309-21.  
Journal code: E7T. ISSN: 0012-1606.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199904  
EW 19990404  
AB Using **transgenic** embryos, we have identified two distinct CNS  
progenitor cell-specific enhancers, each requiring the cooperation of at  
least two independent regulatory sites, within the second intron of the  
rat **nestin** gene. One enhancer is active throughout the  
developing CNS, while the other is specifically active in the ventral  
midbrain. These experiments demonstrate that neural progenitor cells in  
the midbrain constitute a unique subpopulation based upon their ability to  
activate the midbrain regulatory element. Our finding of differential  
enhancer activity from a gene encoding a structural protein reveals a  
previously unrecognized diversity in neural progenitor cell populations.  
Copyright 1999 Academic Press.